arXiv:0905.3506v1 [hep-ph] 21 May 2009

Doubly Heavy Baryon Zero-Recoil Semileptonic Decay in Heavy Quark-Diquark Symmetry

Jie Hu*[1]

[1] Department of Physics, Duke University,
Durham, North Carolina 27708, USA

(Dated: May 21, 2009)

Abstract

This paper studies the leading chiral corrections to heavy quark-diquark symmetry predictions for doubly heavy baryon semileptonic decay form factors. We derive the couplings of heavy diquarks to weak currents in the limit of heavy quark-diquark symmetry, and construct the chiral Lagrangian for doubly heavy baryons coupled to weak currents. We calculate chiral corrections to doubly heavy baryon zero-recoil semileptonic decay for both unquenched and partially quenched QCD. This theory is used to derive chiral extrapolation formulae for measurements of the doubly heavy baryon zero-recoil semileptonic decay form factors in lattice QCD simulations.

---

* Electronic address: hujie@phy.duke.edu

I. INTRODUCTION

Motivation for this work comes from the SELEX experiment's recent observation of states which have been tentatively interpreted as doubly charm baryons [1, 2, 3]. While the masses and the hyperfine splittings of the observed states are consistent with theoretical predictions from some quark models [4] and quenched lattice QCD [5, 6, 7], many other aspects are difficult to understand theoretically. Therefore model-independent predictions for doubly heavy baryon properties would be helpful. Chiral and heavy quark symmetries are useful approximate symmetries of QCD for predicting low-energy properties of hadrons containing a single heavy quark. Heavy quark-diquark symmetry relates heavy mesons to doubly heavy antibaryons. The heavy quark-diquark symmetry prediction for the doubly charm hyperfine splitting [4, 5, 13, 14] is within 25% $\sim$ 30% of the quenched lattice QCD calculation. More observables are needed to see if heavy quark-diquark symmetry can be applied to charm and bottom. In this paper we will study the leading chiral corrections to heavy quark-diquark symmetry predictions for zero-recoil doubly heavy baryon semileptonic decay form factors.

Heavy quark-diquark symmetry relates mesons with a single heavy quark ($Q\bar{q}$) to antibaryons with two heavy antiquarks ($\bar{Q}\bar{Q}\bar{q}$). Savage and Wise [8] wrote an effective Lagrangian for heavy quarks and diquarks using heavy quark effective theory (HQET) [9] and obtained a prediction for doubly heavy baryon hyperfine splittings. HQET considers only the scales Λ_{QCD} and m_Q and is formulated as an expansion in Λ_{QCD}/m_Q. Hadrons with two or more heavy quarks, such as quarkonia or doubly heavy baryons, are characterized by the additional scales $m_Q v$ and $m_Q v^2$, where v is the typical velocity of the heavy quark within the bound state. The correct theory of hadrons with two or more heavy quarks is Non-Relativistic Quantum Chromodynamics (NRQCD) [10, 11, 12]. Refs. [13, 14] derived effective Lagrangians for heavy diquarks within the framework of NRQCD and corrected a factor of 2 in the prediction for hyperfine splittings of Ref. [8]. In this paper we will use these techniques to derive the couplings of heavy diquarks to weak currents.

For studying the low energy dynamics of hadrons containing diquarks, it is useful to build an effective theory which incorporates the relevant symmetries of QCD. Heavy hadron chiral perturbation theory (HHχPT) [15, 16, 17] has heavy hadrons, Goldstone bosons, and photons as its elementary degrees of freedom and incorporates chiral and heavy quark symmetries. Ref. [18] derived an effective Lagrangian for doubly heavy baryons incorporating heavy quark-diquark symmetry and used this theory to study low energy strong and electromagnetic interactions of doubly heavy baryons. In this paper, we will apply this theory to calculate chiral corrections to semileptonic decay at zero-recoil.

Lattice gauge theory is a useful tool to address the nonperturbative dynamics of quarks and gluons. First principles numerical calculations of doubly heavy baryons from lattice QCD can be compared with experiment to test our understanding of QCD. Lattice calculation of decay matrix elements of heavy mesons and doubly heavy baryons can help determine the reliability of heavy quark-diquark symmetry for charm and bottom hadrons. HPQCD and MILC collaborations are planning to calculate semileptonic decay form factors of heavy mesons in the near future. In lattice QCD, they do simulations with unphysical sea quark masses which need to be extrapolated to the physical values. Chiral Lagrangian in quenched and partially quenched theory is useful for the chiral extrapolations of lattice data. In this paper, we will extend the chiral Lagrangian with heavy quark-diquark symmetry to include effects of quenching and partially quenching, and use it to derive formulae for the chiral extrapolation of zero-recoil semileptonic decay of double heavy baryons in lattice QCD simulations.

The rest of this paper is organized as follows. In Sec. II, we use NRQCD to derive the couplings of heavy diquarks to weak currents consistent with heavy quark-diquark symmetry. In Sec. III, we construct the chiral Lagrangian for doubly heavy baryons coupled to weak currents and calculate the doubly heavy baryon zero-recoil semileptonic decay matrix elements. We calculate the chiral corrections to doubly heavy baryon zero-recoil semileptonic decay using χPT. In heavy quark-diquark limit the corrections vanish. In Sec. IV, we extend the chiral Lagrangian with heavy quark-diquark symmetry to the partially quenched theory and derive formulae for chiral extrapolation of doubly heavy baryon zero-recoil semileptonic decay form factors in lattice QCD simulations. We compare chiral corrections to the zero-recoil doubly heavy baryon semileptonic decay form factor in PQχPT with the corresponding corrections in χPT. In Sec. V, a brief summary is given. Some useful formulae are collected in the Appendix.

II. COUPLINGS OF HEAVY DIQUARK TO WEAK CURRENTS FROM VNRQCD

NRQCD is the nonrelativistic effective theory which describes the dynamics of nonrelativistic heavy quarks. In NRQCD there are four important energy scales, m_Q, the typical momentum of the heavy quarks within the bound state $m_Q v$, the typical kinetic energy of the heavy quarks $m_Q v^2$ and Λ_{QCD}. The total momentum of the heavy quark field is taken to be the sum of the label momentum $\mathbf{p}$ and the residual momentum $\mathbf{k}$ as $\mathbf{p}_{total} = \mathbf{p}(\sim m_Q v) + \mathbf{k}(\sim m_Q v^2)$. vNRQCD is an effective theory for NRQCD which has a consistent v expansion constructed by Luke, Manohar and Rothstein [12]. The leading order vNRQCD Lagrangian to $O(v^5)$ for the $\bar{Q}\bar{Q}$ sector for $\bar{b}, \bar{c}$ antiquarks is

$$\begin{aligned}\mathcal{L} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} &+ \sum_{f=b,c}\sum_{\mathbf{p}} \chi_{\mathbf{p}}^{f\dagger}\left(iD_0 - \frac{(\mathbf{p}-i\mathbf{D})^2}{2m_{Q_f}} + \frac{g_s}{2m_{Q_f}}\boldsymbol{\sigma}\cdot\mathbf{B}\right)\chi_{\mathbf{p}}^{f} \\ &- \frac{1}{2}\sum_{f=b,c}\sum_{\mathbf{p},\mathbf{q}} \frac{g_s^2}{(\mathbf{p}-\mathbf{q})^2}\chi_{\mathbf{q}}^{f\dagger}\bar{T}^A\chi_{\mathbf{p}}^{f}\chi_{-\mathbf{q}}^{f\dagger}\bar{T}^A\chi_{-\mathbf{p}}^{f} - \sum_{\mathbf{p},\mathbf{q}}\frac{g_s^2}{(\mathbf{p}-\mathbf{q})^2}\chi_{\mathbf{q}}^{b\dagger}\bar{T}^A\chi_{\mathbf{p}}^{b}\chi_{-\mathbf{q}}^{c\dagger}\bar{T}^A\chi_{-\mathbf{p}}^{c} + \cdots,\end{aligned} \tag{1}$$

where the ellipsis represents high order terms. $\chi_{\mathbf{p}}^{f}$ is a nonrelativistic antiquark field with flavor f which annihilates an antiquark. The kinetic energy D_0 and momentum $\mathbf{p}$ of the heavy quarks are $O(m_Q v^2)$ and $O(m_Q v)$, respectively. $\bar{T}^A$ is $SU(3)$ color generators for $\bar{3}$ representation, $\mathbf{B}$ is chromomagnetic field, D_0 and $\mathbf{D}$ are the time and spatial components of the gauge covariant derivative. In order to derive an effective Lagrangian for diquarks we follow the methods of Ref. [14]. We use the spin and color Fierz identities

$$\delta_{\alpha\delta}\delta_{\beta\gamma} = -\frac{1}{2}(\sigma^i\epsilon)_{\alpha\beta}(\epsilon\sigma^i)_{\gamma\delta} + \frac{1}{2}\epsilon_{\alpha\beta}\epsilon^T_{\gamma\delta}\,, \tag{2}$$

$$\bar{T}^A_{il}\bar{T}^A_{jk} = \frac{2}{3}\sum_m \frac{1}{2}\epsilon_{mij}\epsilon_{mkl} + \frac{1}{3}\sum_{(mn)} d^{(mn)}_{ij}d^{(mn)}_{kl}, \tag{3}$$

to project the potential onto the $\mathbf{3}$ and $\bar{\mathbf{6}}$ channels and to decompose the antiquark bilinears such as $\chi_{\mathbf{p}}^{f}\chi_{-\mathbf{p}}^{f}$ into operators with spin 0 and spin 1. In Eq.(3) The Greek letters denote spin indices, the σ^i denote the Pauli matrices and $\epsilon = i\sigma^2$. In Eq.(3), the Roman letters denote the color indices and the matrices $d^{(mn)}_{ij}$ are symmetric matrices in color space defined by

$$d^{(mn)}_{ij} = \begin{cases} (\delta^m_i\delta^n_j + \delta^n_i\delta^m_j)/\sqrt{2} & m \neq n \\ \delta^m_i\delta^n_j & m = n \end{cases}. \tag{4}$$

For bb and cc, the diquark state must be in either$(\mathbf{3})_C(\mathbf{3})_S$ or $(\bar{\mathbf{6}})_C(\mathbf{1})_S$ by the Pauli principle. However, there is no restriction on the color and spin for mixed flavor. After a Fourier transform to obtain position space potentials, the Lagrangian can be rewritten as

$$\begin{aligned}
\mathcal{L} \; = \; & -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} + \sum_{f=b,c}\sum_{\mathbf{p}} \chi_{\mathbf{p}}^{f\dagger}\left(iD_0 - \frac{(\mathbf{p}-i\mathbf{D})^2}{2m_{Q_f}} + \frac{g_s}{2m_{Q_f}}\boldsymbol{\sigma}\cdot\mathbf{B}\right)\chi_{\mathbf{p}}^{f} \\
& - \frac{1}{2}\sum_{f=b,c}\int d^3\mathbf{r}V^{(3)}(r)\left(\sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}}\frac{\epsilon_{mij}}{2}(\chi_{\mathbf{q}}^{f\dagger})_i\boldsymbol{\sigma}\epsilon(\chi_{-\mathbf{q}}^{f\dagger})_j\right)\cdot\left(\sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}}\frac{\epsilon_{mkl}}{2}(\chi_{-\mathbf{p}}^{f})_k\epsilon\boldsymbol{\sigma}(\chi_{\mathbf{p}}^{f})_l\right) \\
& - \int d^3\mathbf{r}V^{(3)}(r)\left(\sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}}\frac{\epsilon_{mij}}{2}(\chi_{\mathbf{q}}^{b\dagger})_i\boldsymbol{\sigma}\epsilon(\chi_{-\mathbf{q}}^{c\dagger})_j\right)\cdot\left(\sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}}\frac{\epsilon_{mkl}}{2}(\chi_{-\mathbf{p}}^{c})_k\epsilon\boldsymbol{\sigma}(\chi_{\mathbf{p}}^{b})_l\right) \\
& - \int d^3\mathbf{r}V^{(3)}(r)\left(-\sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}}\frac{\epsilon_{mij}}{2}(\chi_{\mathbf{q}}^{b\dagger})_i\epsilon(\chi_{-\mathbf{q}}^{c\dagger})_j\right)\left(\sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}}\frac{\epsilon_{mkl}}{2}(\chi_{-\mathbf{p}}^{c})_k\epsilon^T(\chi_{\mathbf{p}}^{b})_l\right) \\
& + \cdots ,
\end{aligned} \tag{5}$$

where the ellipsis reflexes to terms in which the diquark is in the $\bar{6}$ of color, $V^{(3)}(r) = -\frac{2}{3}\frac{\alpha_s}{r}$ and $V^{(\bar{6})}(r) = \frac{1}{3}\frac{\alpha_s}{r}$. The first line includes the gauge boson fields, the kinetic terms for the antiquarks and the leading spin symmetry breaking interactions which generate hyperfine splittings. The next three lines are the quartic terms of diquark fields. In order to get the interaction for doubly heavy baryon semileptonic decay, we need the Lagrangian for diquarks of different flavor. The diquark fields for $\bar{c}\bar{c}$ and $\bar{b}\bar{b}$ can be introduced using the Hubbard-Strantonovich transformation as in Ref. [14]. We add to the Lagrangian:

$$\begin{aligned}
\Delta\mathcal{L} \; = \; & \frac{1}{2}\sum_{f=b,c}\int d^3\mathbf{r}V^{(3)}(r)\left(\mathbf{T}_{\mathbf{r}}^{m\dagger} - \sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}}\frac{\epsilon_{mij}}{2}(\chi_{\mathbf{q}}^{f\dagger})_i\boldsymbol{\sigma}\epsilon(\chi_{-\mathbf{q}}^{f\dagger})_j\right) \\
& \cdot\left(\mathbf{T}_{\mathbf{r}}^{m} - \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}}\frac{\epsilon_{mkl}}{2}(\chi_{-\mathbf{p}}^{f})_k\epsilon\boldsymbol{\sigma}(\chi_{\mathbf{p}}^{f})_l\right) \\
& + \int d^3\mathbf{r}V^{(3)}(r)\left(\tilde{\mathbf{T}}_{\mathbf{r}}^{m\dagger} - \sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}}\frac{\epsilon_{mij}}{2}(\chi_{\mathbf{q}}^{b\dagger})_i\boldsymbol{\sigma}\epsilon(\chi_{-\mathbf{q}}^{c\dagger})_j\right) \\
& \cdot\left(\tilde{\mathbf{T}}_{\mathbf{r}}^{m} - \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}}\frac{\epsilon_{mkl}}{2}(\chi_{-\mathbf{p}}^{c})_k\epsilon\boldsymbol{\sigma}(\chi_{\mathbf{p}}^{b})_l\right) \\
& + \int d^3\mathbf{r}V^{(3)}(r)\left(T_{\mathbf{r}}^{\prime m\dagger} + \sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}}\frac{\epsilon_{mij}}{2}(\chi_{\mathbf{q}}^{b\dagger})_i\epsilon(\chi_{-\mathbf{q}}^{c\dagger})_j\right) \\
& \times\left(T_{\mathbf{r}}^{\prime m} - \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}}\frac{\epsilon_{mkl}}{2}(\chi_{-\mathbf{p}}^{c})_k\epsilon^T(\chi_{\mathbf{p}}^{b})_l\right),
\end{aligned} \tag{6}$$

where $m = 1,2,3$ is the color index. The $\mathbf{T}^m = \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}} \frac{\epsilon_{mkl}}{2} (\chi^f_{-\mathbf{p}})_k \epsilon \boldsymbol{\sigma} (\chi^f_{\mathbf{p}})_l$ is the spin-1 $\bar{f}\bar{f}$ field, $\tilde{\mathbf{T}}^m = \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}} \frac{\epsilon_{mkl}}{2} (\chi^c_{-\mathbf{p}})_k \epsilon \boldsymbol{\sigma} (\chi^b_{\mathbf{p}})_l$ is the spin-1 $\bar{b}\bar{c}$ field, and the $T'^m = \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}} \frac{\epsilon_{mkl}}{2} (\chi^c_{-\mathbf{p}})_k \epsilon^T (\chi^b_{\mathbf{p}})_l$ is the spin-0 $\bar{b}\bar{c}$ field. The quartic terms of antiquark fields cancel and we are left with the following action

$$\begin{aligned}
\mathcal{L} =& \\
\tfrac{1}{2} \sum_{f=b,c} \int & d^3\mathbf{r} V^{(3)}(r) \left(\mathbf{T}^{m\dagger}_{\mathbf{r}} \mathbf{T}^m_{\mathbf{r}} - \mathbf{T}^{m\dagger}_{\mathbf{r}} \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}} \frac{\epsilon_{mkl}}{2} (\chi^f_{-\mathbf{p}})_k \epsilon \boldsymbol{\sigma} (\chi^f_{\mathbf{p}})_l - \mathbf{T}^m_{\mathbf{r}} \sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{\epsilon_{mij}}{2} (\chi^{f\dagger}_{\mathbf{q}})_i \boldsymbol{\sigma} \epsilon (\chi^{f\dagger}_{-\mathbf{q}})_j \right) \\
+ \int & d^3\mathbf{r} V^{(3)}(r) \left(\tilde{\mathbf{T}}^{m\dagger}_{\mathbf{r}} \tilde{\mathbf{T}}^m_{\mathbf{r}} - \tilde{\mathbf{T}}^{m\dagger}_{\mathbf{r}} \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}} \frac{\epsilon_{mkl}}{2} (\chi^c_{-\mathbf{p}})_k \epsilon \boldsymbol{\sigma} (\chi^b_{\mathbf{p}})_l - \tilde{\mathbf{T}}^m_{\mathbf{r}} \sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{\epsilon_{mij}}{2} (\chi^{b\dagger}_{\mathbf{q}})_i \boldsymbol{\sigma} \epsilon (\chi^{c\dagger}_{-\mathbf{q}})_j \right) \\
+ \int & d^3\mathbf{r} V^{(3)}(r) \left(T'^{m\dagger}_{\mathbf{r}} T'^m_{\mathbf{r}} - T'^{m\dagger}_{\mathbf{r}} \sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{r}} \frac{\epsilon_{mkl}}{2} (\chi^c_{-\mathbf{p}})_k \epsilon^T (\chi^b_{\mathbf{p}})_l + T'^m_{\mathbf{r}} \sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{\epsilon_{mij}}{2} (\chi^{b\dagger}_{\mathbf{q}})_i \epsilon (\chi^{c\dagger}_{-\mathbf{q}})_j \right) . \qquad (7)
\end{aligned}$$

The Feynman rules for the couplings of the diquark fields to the pairs of antiquarks from the interaction terms in Eq.(7) are shown in Fig. 1. Integrating out the fields $\mathbf{T}^m_{\mathbf{r}}$, $\tilde{\mathbf{T}}^m_{\mathbf{r}}$ and $T'^m_{\mathbf{r}}$ will give us back the original NRQCD Lagrangian. Integrating out the antiquark fields $\chi^b_{\mathbf{p}}$ and $\chi^c_{-\mathbf{p}}$ will yield an effective action for the diquark fields $\mathbf{T}^m_{\mathbf{r}}$, $\tilde{\mathbf{T}}^m_{\mathbf{r}}$ and $T'^m_{\mathbf{r}}$. The $\bar{b}\bar{c}$ fields kinetic terms are the same as those of $\bar{c}\bar{c}$ fields in Ref. [14], with the exception that the diquark reduced mass is now

$$\frac{m_Q}{2} \to \mu_Q = \frac{m_b + m_c}{m_b m_c} \, . \qquad (8)$$

The coupling of the diquarks to weak currents is obtained by evaluating graphs in Fig. 2. The flavor changing weak currents are given by $J^\mu_{QCD} = \bar{c}\gamma^\mu(1-\gamma_5)b$. By matching onto NRQCD, the weak current is rewritten by $J^\mu_{NR} = \chi^{c\dagger}(\delta^{\mu 0} - \delta^{\mu i}\sigma^i)\chi^b + \cdots$ where the ellipsis reflexes to higher order terms. For the lowest order weak currents from Fig. 2 we find the couplings of diquark to weak currents. The Lagrangian for the weak interactions of diquarks


$\mathbf{p}, i, \alpha$
$\mathbf{T}^m_{\mathbf{r}}, \tilde{\mathbf{T}}^m_{\mathbf{r}}$
$-ie^{-i\mathbf{p}\cdot\mathbf{r}}V^{(3)}(r)\frac{\epsilon_{mij}}{2}(\boldsymbol{\sigma}\epsilon)_{\alpha\beta}$
$-\mathbf{p}, j, \beta$
$\mathbf{p}, i, \alpha$
$T'^m_{\mathbf{r}}$
$ie^{-i\mathbf{p}\cdot\mathbf{r}}V^{(3)}(r)\frac{\epsilon_{mij}}{2}\epsilon_{\alpha\beta}$
$-\mathbf{p}, j, \beta$

FIG. 1: Feynman rules for the coupling of the diquarks $\mathbf{T}^m_{\mathbf{r}}$, $\tilde{\mathbf{T}}^m_{\mathbf{r}}$ and $T'^m_{\mathbf{r}}$ to antiquarks. The Greek letters denote spin indices and the Roman letters refer to the color indices.



FIG. 2: One loop diagram contributing to the coupling of composite anti-diquark fields and weak currents.

is

$$\begin{aligned}
\mathcal{L} &= J^\mu_{QQ}(J_W)_\mu \\
&= \int d^3\mathbf{r}\left(-\delta^{\mu 0}\mathbf{T}^{i\dagger}_{\mathbf{r}}\cdot\tilde{\mathbf{T}}^i_{\mathbf{r}} - i\delta^{\mu n}(\mathbf{T}^{i\dagger}_{\mathbf{r}}\times\tilde{\mathbf{T}}^i_{\mathbf{r}})^n - \delta^{\mu n}(T^{i\dagger}_{\mathbf{r}})^n T'^i_{\mathbf{r}}\right)(J_W)_\mu \\
&= \eta\left(-\delta^{\mu 0}\mathbf{T}^{i\dagger}\cdot\tilde{\mathbf{T}}^i - i\delta^{\mu n}(\mathbf{T}^{i\dagger}\times\tilde{\mathbf{T}}^i)^n - \delta^{\mu n}(T^{i\dagger})^n T'^i\right)(J_W)_\mu + \cdots , \qquad (9)
\end{aligned}$$

where J_W is the weak current. In the last line we expand the diquark fields to the lowest order and write the lagrangian in terms of local currents. The $\cdots$ denotes the excited diquark states contribution and the factor η can be interpreted as the spatial wavefunction overlap of the ground state initial diquark with the ground state final diquark system, which is not predicted by symmetry, i.e. $\eta = \int d^3\mathbf{r}\phi^*_{bc}(\mathbf{r})\phi_{cc}(\mathbf{r})\, e^{i\mathbf{q}\cdot\mathbf{r}}$. Our results differ from Ref. [21] by a different relative sign of the coupling to the zeroth component of the weak current and that of the spatial component of the weak current.

III. DOUBLY HEAVY BARYON SEMILEPTONIC DECAY

In the heavy quark limit, the lowest mass states of the spin-0 heavy meson, P, and the spin-1 heavy meson, P^*, are degenerate, and therefore combined into a single field, H_a,

$$H_a = \vec{P}^*_a \cdot \vec{\sigma} + P_a \,, \tag{10}$$

where a is an $SU(3)$ flavor anti-fundamental index and the $\vec{\sigma}$ are the Pauli matrices. Similarly, the ground state degenerate doubly heavy baryon doublet, $T_{i\beta}$ [1], consists of a spin-1/2 doubly heavy baryon, $\Xi_{a,\gamma}$, and a spin-3/2 doubly heavy baryon, $\Xi^*_{a,i\beta}$,

$$T_{a,i\beta} = \sqrt{2}\left(\Xi^*_{a,i\beta} + \frac{1}{\sqrt{3}}\Xi_{a,\gamma}\,\sigma^i_{\gamma\beta}\right)\,, \tag{11}$$

where $\Xi^*_{a,i\beta}$ and $\Xi_{a,\gamma}$ are the spin-$\frac{3}{2}$ and spin-$\frac{1}{2}$ fields, respectively. The $i = 1, 2, 3$ are diquark spin indices and $\beta = 1, 2$ are light antiquark spin indices. The field $\Xi^*_{a,i\beta}$ obeys the constraint $\Xi^*_{a,i\beta}\,\sigma^i_{\beta\gamma} = 0$. Heavy quark-diquark symmetry relates the heavy mesons to doubly heavy antibaryons. The effective Lagrangian for doubly heavy baryon and heavy meson ground state doublets in the heavy hadron rest frame was constructed in Ref. [18]

$$\mathcal{L} = \mathrm{Tr}[\mathcal{H}^\dagger_a (iD_0)_{ba}\mathcal{H}_b] - g\,\mathrm{Tr}[\mathcal{H}^\dagger_a \mathcal{H}_b\,\vec{\sigma}\cdot\vec{A}_{ba}] + \frac{\Delta_H}{4}\mathrm{Tr}[\mathcal{H}^\dagger_a\,\Sigma^i\,\mathcal{H}_a\,\sigma^i]\,, \tag{12}$$

where $\mathcal{H}_{a,\mu\beta} = H_{a,\alpha\beta} + T_{a,i\beta}$ is a 5×2 matrix field which transforms as tensor products of the five component field $\mathcal{Q}_\mu$ (which corresponding to the two heavy quark spin states and the three diquark spin states) and a two-component light antiquark spinor. Here the index

---

[1] In this section, the T is the doubly heavy baryon field, although T is diquark field in the previous section.

μ takes on values between 1 and 5, $\alpha, \beta = 1$ or 2, and $i = 3, 4$, or 5. The first term is the kinetic term, the second term is the coupling to axial current and these two terms respect the $SU(5)$ heavy quark-diquark symmetry. The third term is the hyperfine splitting term which is the leading heavy quark symmetry breaking. The $\vec{\Sigma}_{\mu\nu}$ are the 5×5 matrices

$$\vec{\Sigma}_{\mu\nu} = \begin{pmatrix} \vec{\sigma}_{\alpha\beta} & 0 \\ 0 & \vec{\mathcal{T}}_{jk} \end{pmatrix} , \tag{13}$$

and $(\mathcal{T}^i)_{jk} = -i\epsilon_{ijk}$. Δ_H is the hyperfine splitting of the charmed mesons which is related to the hyperfine fplitting of the doubly charm baryon by heavy quark-diquark symmetry. $\vec{A}_{ba}$ is the spatial part of the axial vector current.

The relevant terms of Eq. (12) for our calculation are

$$\begin{aligned} \mathcal{L} &= \tilde{T}^{+}_{\beta i} iD_0 \tilde{T}_{i\beta} - g\tilde{T}^{+}_{\beta i}\tilde{T}_{i\beta'}\vec{\sigma}_{\beta'\beta}\cdot\vec{A} + \frac{\Delta_H}{4}\tilde{T}^{+}_{\beta j}(-i\epsilon_{ijk})\tilde{T}_{k\beta'}\sigma^{i}_{\beta'\beta} \\ &+ T^{+}_{\beta i} iD_0 T_{i\beta} - gT^{+}_{\beta i}T_{i\beta'}\vec{\sigma}_{\beta'\beta}\cdot\vec{A} + \left(\frac{m_c/2}{\mu_Q}\right)\frac{\Delta_H}{4}T^{+}_{\beta j}(-i\epsilon_{ijk})T_{k\beta'}\sigma^{i}_{\beta'\beta} \\ &+ T'^{+}_{\beta} iD_0 T'_{\beta} - g'T'^{+}_{\beta}T'_{\beta'}\vec{\sigma}_{\beta'\beta}\cdot\vec{A} . \end{aligned} \tag{14}$$

The composite fields are decomposed as

$$\begin{aligned} (T_{\bar{c}\bar{c}\bar{q}})_{i\beta} &= \sqrt{2}\left(\Xi^{*}_{\bar{c}\bar{c}\bar{q},i\beta} + \frac{1}{\sqrt{3}}\Xi_{\bar{c}\bar{c}\bar{q},\alpha}\sigma^{i}_{\alpha\beta}\right) , \\ (\tilde{T}_{\bar{b}\bar{c}\bar{q}})_{i\beta} &= \sqrt{2}\left(\Xi^{*}_{\bar{b}\bar{c}\bar{q},i\beta} + \frac{1}{\sqrt{3}}\Xi_{\bar{b}\bar{c}\bar{q},\alpha}\sigma^{i}_{\alpha\beta}\right) , \\ (T'_{\bar{b}\bar{c}\bar{q}})_{\beta} &= \sqrt{2}\,\Xi'_{\bar{b}\bar{c}\bar{q},\beta} , \end{aligned} \tag{15}$$

where $\Xi^{*}_{i\beta}$ and Ξ_{α} are the spin-$\frac{3}{2}$ and spin-$\frac{1}{2}$ fields, respectively. Here, we have suppressed the flavor indices. $T_{i\beta}$ and $\tilde{T}_{i\beta}$ are the ground state doublet of $\bar{c}\bar{c}\bar{q}$ and $\bar{b}\bar{c}\bar{q}$, respectively, and the diquarks are in spin-1 channel. T'_{β} is the spin-$\frac{1}{2}$ ground state of $\bar{b}\bar{c}\bar{q}$ where the diquark is in spin-0 channel. The hyperfine splitting of the doubly heavy baryons are related to that of the charmed mesons by heavy quark-diquark symmetry, $m_{\Xi^{*}_{cc}} - m_{\Xi_{cc}} = \frac{3}{4}\Delta_H$ and $m_{\Xi^{*}_{bc}} - m_{\Xi_{bc}} = \left(\frac{m_c/2}{\mu_Q}\right)\frac{3}{4}\Delta_H$ where the μ_Q is the reduced mass of diquark bc.

Eq.14 describes the low energy strong and electromagnetic interactions in the doubly heavy baryon rest frame in which the doubly heavy baryon four-velocity is conserved (up to $O(\Lambda_{QCD}/m_Q)$ corrections). For a process such as the weak decay, in which the initial and final baryons have different four-velocities, the covariant representation of baryon fields is needed. However, for studying the zero-recoil semileptonic decay, in which the doubly heavy baryon four-velocity is conserved, it is possible to work in the baryon rest frame. Therefore we are allowed to match the weak currents coupling of diquark onto a current operator in the effective theory for doubly heavy baryons. The Lagrangian for semileptonic decays of doubly heavy baryon is

$$\mathcal{L} \;=\; i\eta\; T^{+}_{\beta i}\epsilon_{ijk}\tilde{T}_{k\beta}J^{k}_{W} - \eta\; T^{+}_{\beta i}\tilde{T}_{i\beta}J^{0}_{W} + i\eta\; T^{+}_{\beta i}T'_{\beta}J^{i}_{W}\;. \tag{16}$$

The couplings of the doubly heavy baryon to the weak currents is obtained by demanding that its diquark index on the doubly heavy baryons couple to the weak currents as in Eq.(9).

Expanding out Eq.(16) in terms of doubly heavy baryon fields, we find the following weak current matrix elements in terms of the initial and final spinor.

$$\langle \Xi_{\bar{c}\bar{c}\bar{q}}|J^{\mu}_{W}|\Xi_{\bar{b}\bar{c}\bar{q}}\rangle = \eta\,\bar{u}_{\alpha'}\left(-2i(1+\delta_1)\,\delta_{\alpha\alpha'}\delta^{\mu 0} - \frac{4i}{3}(1+\delta_2)\,\sigma^{j}_{\alpha\alpha'}\delta^{\mu j}\right)u_{\alpha}\,, \tag{17}$$

$$\langle \Xi^{*}_{\bar{c}\bar{c}\bar{q}}|J^{\mu}_{W}|\Xi_{\bar{b}\bar{c}\bar{q}}\rangle = \eta\,\bar{u}_{i\beta}\left(\frac{2i}{\sqrt{3}}(1+\delta_3)\,\delta_{\alpha\beta}\delta^{\mu i}\right)u_{\alpha}\,, \tag{18}$$

$$\langle \Xi_{\bar{c}\bar{c}\bar{q}}|J^{\mu}_{W}|\Xi^{*}_{\bar{b}\bar{c}\bar{q}}\rangle = \eta\,\bar{u}_{\alpha}\left(\frac{2i}{\sqrt{3}}(1+\delta_4)\,\delta_{\alpha\beta}\delta^{\mu i}\right)u_{i\beta}\,, \tag{19}$$

$$\langle \Xi^{*}_{\bar{c}\bar{c}\bar{q}}|J^{\mu}_{W}|\Xi^{*}_{\bar{b}\bar{c}\bar{q}}\rangle = \eta\,\bar{u}_{k\beta}\left(-2i(1+\delta_5)\,\delta_{\alpha\beta}\delta_{ik}\delta^{\mu 0} + 2i(1+\delta_6)\,\sigma^{j}_{\alpha\beta}\delta_{ik}\delta^{\mu j}\right)u_{i\alpha}\,, \tag{20}$$

$$\langle \Xi_{\bar{c}\bar{c}\bar{q}}|J^{\mu}_{W}|\Xi'_{\bar{b}\bar{c}\bar{q}}\rangle = \eta\,\bar{u}_{\alpha'}\left(-\frac{2}{\sqrt{3}}(1+\delta'_1)\,\sigma^{j}_{\alpha\alpha'}\delta^{\mu j}\right)u'_{\alpha}\,, \tag{21}$$

$$\langle \Xi^{*}_{\bar{c}\bar{c}\bar{q}}|J^{\mu}_{W}|\Xi'_{\bar{b}\bar{c}\bar{q}}\rangle = \eta\,\bar{u}_{i\beta}\left(-2(1+\delta'_2)\,\delta_{\alpha\beta}\delta^{\mu i}\right)u'_{\alpha}\,, \tag{22}$$



FIG. 3: One-loop contributions to the doubly heavy baryon semileptonic decay.

where $(u_\alpha, u'_\alpha, u_{i\beta})$ and $(\bar{u}_\alpha, \bar{u}_{i\beta}, \bar{u}'_\alpha)$ are nonrelativistic spinor for initial and final state, respectively. These form factors are predicted by heavy quark-diquark symmetry. The tree level zero-recoil doubly heavy baryon semileptonic decay matrix elements agree with Ref. [20] up to an overall minus sign[2]. At tree level, the δ_i are zero and we confirm that the δ_i vanish at any order in the heavy quark limit. They will receive corrections from heavy quark symmetry breaking effects. Obtaining those symmetry breaking corrections that come from chiral loop corrections in Fig. 3 is the goal of this paper (δ'_1 and δ'_2 are not calculated in this paper).

The pion-baryon vertex in Fig. 3 is generated from the pion-baryon interaction terms in Eq.(14) and the weak current vertex is from the expanded weak currents terms in Eq.(16). There are also loops for the wavefunction renormalization which are not shown here. The chiral loop corrections to semileptonic decay form factors are given in the Appendix and a table of δ_i with different values of μ is given in Table I for both $q = u, d$ and $q = s$. The PQχPT corrections will be discussed in the next section. Here we work in the isospin limit, so $m_u = m_d$. We take the the reduced mass of diquark bc to be $\mu_Q = \frac{m_b m_c}{m_b + m_c} \approx m_c$. We

---

[2] They give results for Ξ_{bc} while we give the results for $\Xi_{\bar{b}\bar{c}}$. To obtain their result to the lowest order, we should use the decomposition of the ground state doublet as $T_{i\beta} = \sqrt{2}\left(\Xi^*_{i\beta} - \frac{1}{\sqrt{3}}\sigma^i_{\beta\gamma}\Xi_{a,\gamma}\right)$ and then we agree up-to an overall sign.

	$\delta_1(u,d)$	$\delta_2(u,d)$	$\delta_3(u,d)$	$\delta_4(u,d)$	$\delta_5(u,d)$	$\delta_6(u,d)$
$\mu = 500$ MeV	0.25	0.25	0.07	0.15	-0.05	-0.05
$\mu = 1500$ MeV	0.32	0.32	0.10	0.19	-0.01	-0.02

	$\delta_1(s)$	$\delta_2(s)$	$\delta_3(s)$	$\delta_4(s)$	$\delta_5(s)$	$\delta_6(s)$
$\mu = 500$ MeV	-0.01	-0.01	-0.14	-0.14	-0.16	-0.16
$\mu = 1500$ MeV	-0.0002	-0.001	-0.13	-0.12	-0.13	-0.13

TABLE I: Chiral correstions to doubly heavy baryon containing a u, d or s semileptonic decay form factors for $\mu = 500$ MeV and $\mu = 1500$ MeV.

choose $g = 0.6$ [23] and we set $f_\pi = f_k = f_\eta = 130$ MeV. The experiment value for Δ_H is 140 MeV. In the numerical calculations we use the values of Goldstone masses of $m_\pi \approx 140$ MeV and $m_k \approx 500$ MeV and $m_\eta = \sqrt{\frac{-m_\pi^2+4m_k^2}{3}}$ from the . Then only one parameters can be varied, the renormalization scale μ. In the chiral perturbation theory calculations the logarithmic μ-dependence from loops is cancelled by μ-dependent counterterms we have not included here. We vary μ from 500 MeV to 1500 MeV to obtain an estimate of uncertainty due to the unknown counterterm contributions. We find that, for the doubly heavy baryon containing antiup or antidown quark, the first and second form factors get very big chiral corrections $\sim 25\% - 32\%$ and the fifth and the sixth get small correction $\sim 5\%$. The form factors for doubly heavy baryon containing antistrange quark obtain negative corrections and the first and second form factors are least sensitive to chiral corrections $\sim 1\%$. It would be interesting to see if the observed deviations from heavy quark-diquark symmetry in either experiments or lattice results agree with the chiral corrections.

IV. PQχPT RESULTS

The EFT techniques with heavy quark-diquark symmetry for doubly heavy baryons can also be applied to doubly heavy systems simulated on the lattice. It is useful to extend the χPT results to include lattice artifacts such as quenching and partial quenching. Here we will focus on partially quenched chiral corrections to the doubly heavy baryon zero-recoil semileptonic decay form factors, which provide the formulae needed for chiral extrapolations in lattice calculations. The major modification to the calculation in the previous section is that there is a modified propagator of the Goldstone mesons.

In a lattice calculation, the sea quark masses are often different from valence quark masses. In partially quenched QCD the sea quark masses are different from the valence quark masses, while in quenched QCD,the sea quark contributions are absent. To reproduce the lattice unphysical quarks in field theory, fictitious ghost quarks are added to cancle the valence quark contributions and then we are left with the effects of sea quarks. This is equivalent to the partial quenching artifact in lattice QCD simulations which use different masses for sea and valence quarks. In the limit of $m_{sea} = m_{valence}$, QCD is recovered. PQχPT is useful for lattice extrapolations to physical quark masses. The PQχPT and QχPT pseudoscalar meson sector is described by the Lagrangian [24, 25, 26, 27, 28, 29, 30, 31],

$$\mathcal{L} = \frac{f^2}{8}\text{str}\left(\partial^\mu \Sigma^\dagger \partial_\mu \Sigma\right) + \frac{l}{4}\,\text{str}\left(m_q \Sigma^\dagger + m_q^\dagger \Sigma\right) + \alpha_\Phi \partial^\mu \Phi_0 \partial_\mu \Phi_0 - \mu_0^2 \Phi_0^2, \tag{23}$$

where the field Σ is defined by

$$\Sigma = \exp\left(\frac{2i\Phi}{f}\right) = \xi^2, \tag{24}$$

and the meson fields appear in the $U(6|3)$ matrix,

$$\Phi = \begin{pmatrix} M & \chi^\dagger \\ \chi & \tilde{M} \end{pmatrix}. \tag{25}$$

The operation str() in Eq. (23) is a supertrace over flavor indices, i.e., $\mathrm{str}(A) = \sum_a \epsilon_a A_{aa}$, where $\epsilon_a = 1$ for $a = 1-6$ and $\epsilon_a = -1$ for $a = 7-9$. The M and $\tilde{M}$ matrices contain bosonic mesons, while the χ and $\chi^\dagger$ matrices contain fermionic mesons (one ghost quark with one sea/valence quark).The quark mass matrix is defined as [22]

$$m_q = diag(m_u, m_d, m_s, m_j, m_l, m_r, m_u, m_d, m_s). \tag{26}$$

We work in the isospin limit for both the valence and sea sectors, so $m_u = m_d$ and $m_l = m_j$. We take the strange sea quark to be same as the valence quark, $m_r = m_s$. From the lowest order PQχPT Lagrangian, the meson with quark content $q\bar{q}'$ has the mass of $m^2_{q\bar{q}'} = \frac{\lambda}{f^2}(m_q + m_{q'})$. The $PQ\chi PT$ propagators of the off-diagonal mesons have the usual Klein-Gordon form. The flavor neutral propagator can be conveniently written as [22]

$$\mathcal{G}^{PQ}_{ab} = \epsilon_a \delta_{ab} P_a + \mathcal{P}_{ab}\left(P_a, P_b, P_X\right), \tag{27}$$

where

$$\begin{aligned} P_a &= \frac{i}{q^2 - m^2_{aa} + i\epsilon},\ P_b = \frac{i}{q^2 - m^2_{bb} + i\epsilon},\ P_X = \frac{i}{q^2 - m^2_X + i\epsilon}, \\ \mathcal{P}_{ab}\left(A, B, C\right) &= -\frac{1}{3}\left[\frac{\left(m^2_{aa} - m^2_{jj}\right)\left(m^2_{aa} - m^2_{rr}\right)}{\left(m^2_{aa} - m^2_{bb}\right)\left(m^2_{aa} - m^2_X\right)} A + \frac{\left(m^2_{bb} - m^2_{jj}\right)\left(m^2_{bb} - m^2_{rr}\right)}{\left(m^2_{bb} - m^2_{aa}\right)\left(m^2_{bb} - m^2_X\right)} B \right. \\ &\qquad \left. + \frac{\left(m^2_X - m^2_{jj}\right)\left(m^2_X - m^2_{rr}\right)}{\left(m^2_X - m^2_{aa}\right)\left(m^2_X - m^2_{bb}\right)} C\right]. \end{aligned} \tag{28}$$

where $m^2_X = \frac{1}{3}(m^2_{jj} + 2m^2_{rr})$.

In terms of the 5×2 field, the partially quenched Lagrangian for doubly heavy baryon and heavy meson is [22],

$$\begin{aligned}\mathcal{L}^{PQ} &= \left(\mathcal{H}^\dagger(\mathcal{H}i\overleftarrow{D}_0)\right) - g\left(\mathcal{H}^\dagger\mathcal{H}\cdot\sigma\right) \\ &\quad +\frac{\Delta_H}{4}\left(\mathcal{H}^\dagger\Sigma\cdot\mathcal{H}\sigma\right) + \sigma\left(\mathcal{H}^\dagger\mathcal{H}\mathcal{M}\right) + \sigma'\left(\mathcal{H}^\dagger\mathcal{H}\right)\mathrm{str}(\mathcal{M}),\end{aligned} \tag{29}$$

where $\mathcal{M}$ is the mass operator defined by $\mathcal{M} = \frac{1}{2}\left(\xi m_q \xi + \xi^\dagger m_q \xi^\dagger\right)$ Therefore the baryon mass splittings in PQχPT are given by

$$\begin{aligned}\Delta_{ccqq'} &= m_{\Xi_{\bar c\bar c\bar q'}} - m_{\Xi_{\bar c\bar c\bar q}} = m_{\Xi^*_{\bar c\bar c\bar q'}} - m_{\Xi^*_{\bar c\bar c\bar q}} = -\sigma(m_{q'} - m_q)\,, \\ \Delta_{bcqq'} &= m_{\Xi_{bcq'}} - m_{\Xi_{bcq}} = m_{\Xi^*_{\bar b\bar c\bar q'}} - m_{\Xi^*_{\bar b\bar c\bar q}} = -\sigma(m_{q'} - m_q)\,, \\ \Delta^*_{ccqq'} &= m_{\Xi^*_{\bar c\bar c\bar q'}} - m_{\Xi_{\bar c\bar c\bar q}} = \frac{3}{4}\Delta_H - \sigma(m_{q'} - m_q)\,, \\ \Delta^*_{bcqq'} &= m_{\Xi^*_{\bar b\bar c\bar q'}} - m_{\Xi_{\bar b\bar c\bar q}} = \frac{1}{2}\left(\frac{3}{4}\Delta_H\right) - \sigma(m_{q'} - m_q)\,,\end{aligned} \tag{30}$$

where we take the the reduced mass of diquark bc to be $\mu_Q = \frac{m_b m_c}{m_b + m_c} \approx m_c$, σ is the coupling constant in the mass operator in the Lagrangian in Eq.(29) . We choose $g = 0.6$ [23] and $f = 130$ MeV. The experiment value for Δ_H is 140 MeV. With the values of the $SU(3)$ splitting of the ground state D mesons $m_{D_s} - m_D = -\sigma(m_s - m_u) \approx 100$ MeV, pion mass value $m_\pi^2 = \frac{\lambda}{f^2}(m_u + m_d)$ and kaon mass value $m_k^2 = \frac{\lambda}{f^2}(m_s + m_d)$, we obtain $-\sigma = \frac{100}{m_k^2 - m_\pi^2}\frac{\lambda}{f^2}$. For the mass of η we use the $SU(3)$ prediction, $m_\eta = \sqrt{\frac{-m_\pi^2 + 4m_k^2}{3}}$. Then only three parameters can be varied, the valence pion mass $m_{\pi val} = m_{uu}$, the sea pion mass $m_{\pi sea} = m_{jj}$, and the renormalization scale μ. In the chiral perturbation theory calculations the logarithmic μ-dependence from loops is cancelled by μ-dependent counterterms we have not included here. We vary μ from 500 MeV to 1500 MeV to obtain an estimate of uncertainty due to the unknown counterterm contributions. In the heavy quark limit, the logarithm μ dependence goes away and we obtain the tree level predictions in previous section. Calculating the PQχPT loop diagrams, such as Fig. 3, we find the corrections for the six form factors which we list in Appendix.



FIG. 4: $\Delta\delta_n$ $(q = u)$ as a function of $m_{\pi sea}$ for different values of $m_{\pi val}$. The width of the bands is the results of varying μ between 500 MeV and 1500 MeV.

In the Figs. 4 and 5 we plot the relevant corrections to the form factors $\Delta\delta_i = \delta_{iPQ} - \delta_{i\chi}$ in terms of $m_{\pi sea}$ for different values of $m_{\pi val}$ for $q = u$ and $q = s$. For each value of $m_{\pi val} = 140$ MeV, 280 MeV and 420 MeV, we let the $m_{\pi val}$ range from $m_{\pi val}$ up to the mass of eta-strange, $m_{\eta_s} = m_{ss} \approx 700$ MeV. The bands correspond to varying μ from 500 MeV to 1500 MeV, which is chosen to be the same for both χPT and PQχPT. From the plots, it



FIG. 5: $\Delta\delta_n$ $(q = s)$ as a function of $m_{\pi sea}$ but independent of different values of $m_{\pi val}$. The width of the bands is the results of varying μ between 500 MeV and 1500 MeV.

is easy to see that the partially quenched chiral corrections reproduce the chiral corrections when the sea quark mass goes to the physical valence quark mass. The corrections of the doubly heavy baryon containing a strange quark do not depend on the valence quark masses. As demonstrated by Figs. 4 and 5, the partially quenched chiral non-analytic corrections relevant to the chiral corrections are very insensitive to μ-dependence. The relevant corrections to $\delta_1, \delta_2, \delta_3$ and δ_4 are affected most by sea quark masses $\sim 10\% - 40\%$ and increase with increasing m_{sea} values. The δ_5 and δ_6 are very insensitive to partial quenching effects, receiving $\sim 1\% - 8\%$ small relevant corrections from partial quenching, and are insensitive to sea quark masses, while the other four are not. It would be interesting to test the effects on lattice. Quark-diquark symmetry predictions should be more robust. Deviation from quark-diquark symmetry must be attributed to short distance effects.

V. SUMMARY

In this paper we have used NRQCD to derive the couplings of heavy diquarks to weak currents with heavy quark-diquark symmetry. We constructed the chiral Lagrangian for doubly heavy baryons coupled to weak currents and calculated the tree level predictions for doubly heavy baryon semileptonic weak decay form factors. We calculated the chiral corrections in both unquenched and partially quenched theory. The partially quenched loop corrections to semileptonic decay form factors are given in the Appendix. The formulae will be useful for chiral extrapolation of doubly heavy baryon zero-recoil semileptonic decay form factors in lattice QCD simulations. The chiral corrections of the six form factors range from -15% to 30%. δ_5 and δ_6 for doubly heavy baryon containing antiup or antidown quark receive small corrctions $\sim 5\%$ and δ_1 and δ_2 for doubly heavy baryon containing antistrage quark receive smaller correction $\sim 1\%$. The partially quenched chiral non-analytic corrections relevant to the chiral corrections are very insensitive to μ-dependence. The δ_5 and δ_6 receive small relevant corrections $\sim 1\% - 8\%$ from partial quenching, while the other four are not. It is interesting to see if the size of the corrections in experiments and lattice calculations match the chiral derivations of the calculations.

Acknowledgments

We thank B. Tiburzi for dicussions and especially thank T. Mehen for various discussions and kind guidance. This research is supported in part by DOE grants DE-FG02-05ER41368 and DE-FG02-05ER41376.

APPENDIX: χPT AND PQχPT FORM FACTORS

Here we give the detailed one-loop χPT and PQχPT corrections to zero-recoil semileptonic decay. In the calculation we use $\bar{M S}$ scheme and have not included the counterterms. In the chiral corrections to the six form factors f is pion decay constant, m_i is the mass of the Goldstone boson in the one-loop diagram and C^i_{ab} is a factor which comes from $SU(3)$ Clebsch-Gordan coefficients in the couplings. For loops with charged pions we have $C^{\pi^\pm}_{12} = C^{\pi^\pm}_{21} = 1$, for loops with neutral pions $C^{\pi^0}_{11} = C^{\pi^0}_{22} = \frac{1}{2}$, for loops with kaons $C^K_{3i} = C^K_{i3} = 1$ (i =1 or 2), and for loops with η mesons $C^\eta_{11} = C^\eta_{22} = \frac{1}{6}$ and $C^\eta_{33} = \frac{2}{3}$.

$$\delta_{1\chi}(q) = \sum_{i,q'} C^i_{qq'} \frac{g^2}{(4\pi f)^2} \left(-\frac{1}{9} I(\Delta_{bcqq'}, \Delta_{ccqq'}, m_i, \mu) - \frac{8}{9} I(\Delta^*_{bcqq'}, \Delta^*_{ccqq'}, m_i, \mu) \right) , \quad (31)$$

$$\begin{aligned} \delta_{2\chi}(q) = & \sum_{i,q'} C^i_{qq'} \frac{g^2}{(4\pi f)^2} (\frac{1}{27} I(\Delta_{bcqq'}, \Delta_{ccqq'}, m_i, \mu) - \frac{4}{27} I(\Delta_{bcqq'}, \Delta^*_{ccqq'}, m_i, \mu) \\ & - \frac{4}{27} I(\Delta^*_{bcqq'}, \Delta_{ccqq'}, m_i, \mu) - \frac{20}{27} I(\Delta^*_{bcqq'}, \Delta^*_{ccqq'}, m_i, \mu)) , \end{aligned} \quad (32)$$

$$\begin{aligned} \delta_{3\chi}(q) = & \sum_{i,q'} C^i_{qq'} \frac{g^2}{(4\pi f)^2} (-\frac{8}{27} I(\Delta_{bcqq'}, -\Delta^*_{ccqq'}, m_i, \mu) + \frac{5}{27} I(\Delta_{bcqq'}, \Delta_{ccqq'}, m_i, \mu) \\ & - \frac{4}{27} I(\Delta^*_{bcqq'}, -\Delta^*_{ccqq'}, m_i, \mu) - \frac{20}{27} I(\Delta^*_{bcqq'}, \Delta_{ccqq'}, m_i, \mu)) , \end{aligned} \quad (33)$$

$$\begin{aligned} \delta_{4\chi}(q) = & \sum_{i,q'} C^i_{qq'} \frac{g^2}{(4\pi f)^2} (-\frac{8}{27} I(-\Delta^*_{bcqq'}, \Delta_{ccqq'}, m_i, \mu) + \frac{5}{27} I(-\Delta^*_{bcqq'}, \Delta^*_{ccqq'}, m_i, \mu) \\ & - \frac{4}{27} I(\Delta_{bcqq'}, \Delta_{ccqq'}, m_i, \mu) - \frac{20}{27} I(\Delta_{bcqq'}, \Delta^*_{ccqq'}, m_i, \mu)) , \end{aligned} \quad (34)$$

$$\delta_{5\chi}(q) = \sum_{i,q'} C^i_{qq'} \frac{g^2}{(4\pi f)^2} \left(-\frac{4}{9} I(-\Delta^*_{bcqq'}, -\Delta^*_{ccqq'}, m_i, \mu) - \frac{5}{9} I(\Delta_{bcqq'}, \Delta_{ccqq'}, m_i, \mu) \right) , \quad (35)$$

$$\begin{aligned} \delta_{6\chi}(q) &= \sum_{i,q'} C^i_{qq'} \frac{g^2}{(4\pi f)^2} (-\frac{8}{27} I(-\Delta^*_{bcqq'}, -\Delta^*_{ccqq'}, m_i, \mu) - \frac{4}{27} I(-\Delta^*_{bcqq'}, \Delta_{ccqq'}, m_i, \mu) \\ &- \frac{4}{27} I(\Delta_{bcqq'}, -\Delta^*_{ccqq'}, m_i, \mu) - \frac{11}{27} I(\Delta_{bcqq'}, \Delta_{ccqq'}, m_i, \mu)) \, . \end{aligned} \quad (36)$$

The formulae for χPT corrections to doubly heavy baryon semileptonic decay form factors are given in terms of the following functions,

$$\begin{aligned} I(0,0,m,\mu) &= 0 \, , \\ I(\Delta,\Delta,m,\mu) &= -6\Delta^2 \ln\left(\frac{\mu^2}{m^2}\right) + 4\left(m^2 - 3\Delta^2\right) + 8m\Delta F\left(\frac{\Delta}{m}\right) + 4\left(\Delta^2 - m^2\right) F'\left(\frac{\Delta}{m}\right) , \\ I(\Delta_1,\Delta_2,m,\mu) &= -2\left(\Delta_1^2 + \Delta_1\Delta_2 + \Delta_2^2\right) \ln\left(\frac{\mu^2}{m^2}\right) + 4\left(m^2 - \Delta_1^2 - \Delta_1\Delta_2 - \Delta_2^2\right) \\ &+ \frac{4}{\Delta_2 - \Delta_1} \left((\Delta_2^2 - m^2) F\left(\frac{\Delta_2}{m}\right) m - (\Delta_1^2 - m^2) F\left(\frac{\Delta_1}{m}\right) m \right) , \end{aligned} \quad (37)$$

where

$$F(x) = \begin{cases} -\sqrt{1-x^2}\left(\frac{\pi}{2} - tan^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right)\right) \, , & |x| < 1 \\ \sqrt{x^2-1} \ln\left(x + \sqrt{x^2-1}\right) \, , & |x| \geq 1 \end{cases} , \quad (38)$$

and $F'(x)$ is the first derivative of x.

Next we give the same corrections for the partially quenched case.

$$\begin{aligned} \delta_{1PQ}(q) &= \sum_{q'=j,l,r} \frac{g^2}{(4\pi f)^2} (-\frac{1}{9} I(\Delta_{bcqq'}, \Delta_{ccqq'}, m_{qq'}, \mu) - \frac{8}{9} I(\Delta^*_{bcqq'}, \Delta^*_{ccqq'}, m_{qq'}, \mu)) \\ &+ \frac{g^2}{(4\pi f)^2} (-\frac{1}{9} K(\Delta_{bcqq}, \Delta_{ccqq}, m_{qq}, m_{qq}, \mu) - \frac{8}{9} K(\Delta^*_{bcqq}, \Delta^*_{ccqq}, m_{qq}, m_{qq}, \mu)) \, , \end{aligned} \quad (39)$$

$$\begin{aligned} \delta_{2PQ}(q) &= \sum_{q'=j,l,r} \frac{g^2}{(4\pi f)^2} (\frac{1}{27} I(\Delta_{bcqq'}, \Delta_{ccqq'}, m_{qq'}, \mu) - \frac{4}{27} I(\Delta_{bcqq'}, \Delta^*_{ccqq'}, m_{qq'}, \mu) \\ &- \frac{4}{27} I(\Delta^*_{bcqq'}, \Delta_{ccqq'}, m_{qq'}, \mu) - \frac{20}{27} I(\Delta^*_{bcqq'}, \Delta^*_{ccqq'}, m_{qq'}, \mu)) \\ &+ \frac{g^2}{(4\pi f)^2} (\frac{1}{27} K(\Delta_{bcqq}, \Delta_{ccqq}, m_{qq}, m_{qq}, \mu) - \frac{4}{27} K(\Delta_{bcqq}, \Delta^*_{ccqq}, m_{qq}, m_{qq}, \mu) \\ &- \frac{4}{27} K(\Delta^*_{bcqq}, \Delta_{ccqq}, m_{qq}, m_{qq}, \mu) - \frac{20}{27} K(\Delta^*_{bcqq}, \Delta^*_{ccqq}, m_{qq}, m_{qq}, \mu)) \, , \end{aligned} \quad (40)$$

$$\begin{aligned}
\delta_{3PQ}(q) \;=\; & \sum_{q'=j,l,r}\frac{g^2}{(4\pi f)^2}(-\frac{8}{27}I(\Delta_{bcqq'},-\Delta^*_{ccqq'},m_{qq'},\mu)+\frac{5}{27}I(\Delta_{bcqq'},\Delta_{ccqq'},m_{qq'},\mu)\\
-\; & \frac{4}{27}I(\Delta^*_{bcqq'},-\Delta^*_{ccqq'},m_{qq'},\mu)-\frac{20}{27}I(\Delta^*_{bcqq'},\Delta_{ccqq'},m_{qq'},\mu))\\
+\; & \frac{g^2}{(4\pi f)^2}(-\frac{8}{27}K(\Delta_{bcqq},-\Delta^*_{ccqq},m_{qq},m_{qq},\mu)+\frac{5}{27}K(\Delta_{bcqq},\Delta_{ccqq},m_{qq},m_{qq},\mu)\\
-\; & \frac{4}{27}K(\Delta^*_{bcqq},-\Delta^*_{ccqq},m_{qq},m_{qq},\mu)-\frac{20}{27}K(\Delta^*_{bcqq},\Delta_{ccqq},m_{qq},m_{qq},\mu))\,,
\end{aligned}\tag{41}$$

$$\begin{aligned}
\delta_{4PQ}(q) \;=\; & \sum_{q'=j,l,r}\frac{g^2}{(4\pi f)^2}(-\frac{8}{27}I(-\Delta^*_{bcqq'},\Delta_{ccqq'},m_{qq'},\mu)+\frac{5}{27}I(-\Delta^*_{bcqq'},\Delta^*_{ccqq'},m_{qq'},\mu)\\
-\; & \frac{4}{27}I(\Delta_{bcqq'},\Delta_{ccqq'},m_{qq'},\mu)-\frac{20}{27}I(\Delta_{bcqq'},\Delta^*_{ccqq'},m_{qq'},\mu))\\
+\; & \frac{g^2}{(4\pi f)^2}(-\frac{8}{27}K(-\Delta^*_{bcqq},\Delta_{ccqq},m_{qq},m_{qq},\mu)+\frac{5}{27}K(-\Delta^*_{bcqq},\Delta^*_{ccqq},m_{qq},m_{qq},\mu)\\
-\; & \frac{4}{27}K(\Delta_{bcqq},\Delta_{ccqq},m_{qq},m_{qq},\mu)-\frac{20}{27}K(\Delta_{bcqq},\Delta^*_{ccqq},m_{qq},m_{qq},\mu))\,,
\end{aligned}\tag{42}$$

$$\begin{aligned}
\delta_{5PQ}(q) \;=\; & \sum_{q'=j,l,r}\frac{g^2}{(4\pi f)^2}\left(-\frac{4}{9}I(-\Delta^*_{bcqq'},-\Delta^*_{ccqq'},m_{qq'},\mu)-\frac{5}{9}I(\Delta_{bcqq'},\Delta_{ccqq'},m_{qq'},\mu)\right)\\
+\; & \frac{g^2}{(4\pi f)^2}\left(-\frac{4}{9}K(-\Delta^*_{bcqq},-\Delta^*_{ccqq},m_{qq},m_{qq},\mu)-\frac{5}{9}k(\Delta_{bcqq},\Delta_{ccqq},m_{qq},m_{qq},\mu)\right),
\end{aligned}\tag{43}$$

$$\begin{aligned}
\delta_{6PQ}(q) \;=\; & \sum_{q'=j,l,r}\frac{g^2}{(4\pi f)^2}(-\frac{8}{27}I(-\Delta^*_{bcqq'},-\Delta^*_{ccqq'},m_{qq'},\mu)-\frac{4}{27}I(-\Delta^*_{bcqq'},\Delta_{ccqq'},m_{qq'},\mu)\\
-\; & \frac{4}{27}I(\Delta_{bcqq'},-\Delta^*_{ccqq'},m_{qq'},\mu)-\frac{11}{27}I(\Delta_{bcqq'},\Delta_{ccqq'},m_{qq'},\mu))\\
+\; & \frac{g^2}{(4\pi f)^2}(-\frac{8}{27}K(-\Delta^*_{bcqq},-\Delta^*_{ccqq},m_{qq},m_{qq},\mu)-\frac{4}{27}K(-\Delta^*_{bcqq},\Delta_{ccqq},m_{qq},m_{qq},\mu)\\
-\; & \frac{4}{27}K(\Delta_{bcqq},-\Delta^*_{ccqq},m_{qq},m_{qq},\mu)-\frac{11}{27}K(\Delta_{bcqq},\Delta_{ccqq},m_{qq},m_{qq},\mu))\,,
\end{aligned}\tag{44}$$

where the function $K(\Delta_1,\Delta_2,m,m,\mu)$ which arises from the hairpins is given by

$$K(\Delta_1,\Delta_2,m_a,m_b,\mu)=\mathcal{P}_{ab}\left(I(\Delta_1,\Delta_2,m_a,\mu),I(\Delta_1,\Delta_2,m_b,\mu),I(\Delta_1,\Delta_2,m_X,\mu)\right)\,. \tag{45}$$

---

[1] M. Mattson *et al.* [SELEX Collaboration], Phys. Rev. Lett. **89**, 112001 (2002).

[2] M. A. Moinester *et al.* [SELEX Collaboration], Czech. J. Phys. **53**, B201 (2003).

[3] A. Ocherashvili *et al.* [SELEX Collaboration], Phys. Lett. B **628**, 18 (2005).

[4] D. Ebert, R. N. Faustov, V. O. Galkin and A. P. Martynenko, Phys. Rev. D **66**, 014008 (2002) [arXiv:hep-ph/0201217].

[5] R. Lewis, N. Mathur and R. M. Woloshyn, Phys. Rev. D **64**, 094509 (2001) [arXiv:hep-ph/0107037].

[6] N. Mathur, R. Lewis and R. M. Woloshyn, Phys. Rev. D **66**, 014502 (2002) [arXiv:hep-ph/0203253].

[7] J. M. Flynn, F. Mescia and A. S. B. Tariq [UKQCD Collaboration], JHEP **0307**, 066 (2003) [arXiv:hep-lat/0307025].

[8] M. J. Savage and M. B. Wise, Phys. Lett. B **248**, 177 (1990).

[9] A. V. Manohar and M. B. Wise, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **10**, 1 (2000).

[10] G. T. Bodwin, E. Braaten and G. P. Lepage, Phys. Rev. D **51**, 1125 (1995) [Erratum-ibid. D **55**, 5853 (1997)].

[11] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Nucl. Phys. B **566**, 275 (2000).

[12] M. E. Luke, A. V. Manohar and I. Z. Rothstein, Phys. Rev. D **61**, 074025 (2000).

[13] N. Brambilla, A. Vairo and T. Rösch, Phys. Rev. D **72**, 034021 (2005).

[14] S. Fleming and T. Mehen, Phys. Rev. D **73**, 034502 (2006) [arXiv:hep-ph/0509313].

[15] M. B. Wise, Phys. Rev. D **45**, 2188 (1992).

[16] G. Burdman and J. F. Donoghue, Phys. Lett. B **280**, 287 (1992).

[17] T. M. Yan, H. Y. Cheng, C. Y. Cheung, G. L. Lin, Y. C. Lin and H. L. Yu, Phys. Rev. D **46**, 1148 (1992) [Erratum-ibid. D **55**, 5851 (1997)].

[18] J. Hu and T. Mehen, Phys. Rev. D **73**, 054003 (2006) [arXiv:hep-ph/0511321].

[19] I. W. Stewart, Nucl. Phys. B **529**, 62 (1998) [arXiv:hep-ph/9803227].

[20] J. M. Flynn and J. Nieves, Phys. Rev. D **76**, 017502 (2007) [Erratum-ibid. D **77**, 099901 (2008)] [arXiv:0706.2805 [hep-ph]].

[21] M. J. White and M. J. Savage, Phys. Lett. B **271**, 410 (1991).

[22] T. Mehen and B. C. Tiburzi, Phys. Rev. D **74**, 054505 (2006) [arXiv:hep-lat/0607023].

[23] S. Fajfer and J. F. Kamenik, Phys. Rev. D **74**, 074023 (2006) [arXiv:hep-ph/0606278].

[24] A. Morel, J. Phys. (France) **48**, 1111 (1987).

[25] S. R. Sharpe, Phys. Rev. D **46**, 3146 (1992) [arXiv:hep-lat/9205020].

[26] C. W. Bernard and M. F. L. Golterman, Phys. Rev. D **46**, 853 (1992) [arXiv:hep-lat/9204007].

[27] C. W. Bernard and M. F. L. Golterman, Phys. Rev. D **49**, 486 (1994) [arXiv:hep-lat/9306005].

[28] S. R. Sharpe, Phys. Rev. D **56**, 7052 (1997) [Erratum-ibid. D **62**, 099901 (2000)] [arXiv:hep-lat/9707018].

[29] M. F. L. Golterman and K. C. L. Leung, Phys. Rev. D **57**, 5703 (1998) [arXiv:hep-lat/9711033].

[30] S. R. Sharpe and N. Shoresh, Phys. Rev. D **62**, 094503 (2000) [arXiv:hep-lat/0006017].

[31] S. R. Sharpe and N. Shoresh, Phys. Rev. D **64**, 114510 (2001) [arXiv:hep-lat/0108003].